Table 2. Number, incidence rate ¹, median days away from work ² and relative standard errors ³ of occupational injuries and illnesses involving days away from work ⁴ to selected parts of body with musculoskeletal disorders ⁵ in selected ownerships for Utah, 2008

Ownership	Part of body affected	Total Cases	Incidence Rate	Median Days	Relative Standard Error
private industry	All Selected Parts	2,830	33.8	10	8.4
private industry	2 Trunk	1,710	20.4	10	8.8
private industry	21 Shoulder- including clavicle- scapula	260	3.1	44	13.5
private industry	22 Chest- including ribs- internal organs	40	0.5	15	29.6
private industry	220 Chest- except internal location of diseases or disorders	40	0.5	15	29.6
private industry	23 Back- including spine- spinal cord	1,170	13.9	7	9.3
private industry	230 Back- including spine- spinal cord- unspecified	600	7.1	4	10.6
private industry	231 Lumbar region	480	5.7	10	11.2
private industry	232 Thoracic region	50	0.6	4	26.1
private industry	233 Sacral region	20	0.3	26	36.7
private industry	238 Multiple back regions	20	0.2	89	44.8
private industry	24 Abdomen	190	2.3	14	15.0
private industry	240 Abdomen- except internal location of diseases or disorders	30	0.3	37	34.3
private industry	241 Internal abdominal location- unspecified	20	0.2	15	42.6
private industry	245 Intestines- peritoneum	150	1.8	14	16.6
private industry	2450 Intestines- peritoneum- unspecified	150	1.8	14	16.6
private industry	28 Multiple trunk locations	40	0.4	103	31.1
private industry	3 Upper extremities	800	9.5	10	10.0
private industry	31 Arm(s)	70	0.8	38	23.0
private industry	310 Arm(s)- unspecified	20	0.2	3	40.4
private industry	318 Multiple arm(s) locations	30	0.3	60	36.2
private industry	32 Wrist(s)	630	7.5	10	10.5
private industry	34 Finger(s)- fingernail(s)	80	0.9	10	21.8
private industry	38 Multiple upper extremities locations	20	0.2	11	43.5
private industry	382 Hand(s) and wrist(s)	20	0.2	11	44.9
private industry	4 Lower extremities	290	3.4	18	13.1
private industry	41 Leg(s)	210	2.5	18	14.7
private industry	411 Thigh(s)	20	0.3	2	39.7
private industry	412 Knee(s)	150	1.8	18	16.4
private industry	42 Ankle(s)	70	0.8	40	23.1
private industry	8 Multiple Body Parts	30	0.4	10	33.7
local government	All Selected Parts	170	22.3	15	24.6
		ı			

See footnotes at end of table

Table 2. Number, incidence rate ¹, median days away from work ² and relative standard errors ³ of occupational injuries and illnesses involving days away from work ⁴ to selected parts of body with musculoskeletal disorders ⁵ in selected ownerships for Utah, 2008 -- Continued

Ownership	Part of body affected	Total Cases	Incidence Rate	Median Days	Relative Standard Error
local government local government local government local government local government state government state government	2 Trunk 21 Shoulder- including clavicle- scapula 23 Back- including spine- spinal cord 230 Back- including spine- spinal cord- unspecified 4 Lower extremities 42 Ankle(s) All Selected Parts 2 Trunk	130 40 90 80 20 20 60 40	17.7 5.1 11.9 9.9 3.3 2.9 13.5 8.6	7 97 7 7 29 29 8 3	26.1 40.0 29.4 31.3 48.2 50.8 23.6 25.3
state government state government	23 Back- including spine- spinal cord 231 Lumbar region	30 20	6.9 5.1	3 3	26.5 28.4

 $^{^{1}}$ Incidence rates represent the number of injuries and illnesses per 10,000 full-time workers and were calculated as: (N / EH) X 20,000,000 where,

N = number of injuries and illnesses,

EH = total hours worked by all employees during the calendar year,

20,000,000 = base for 10,000 full-time equivalent workers (working 40 hours per week, 50 weeks per year).

NOTE: Dashes indicate data that do not meet publication guidelines or data for incidence rates less than .05 per 10,000 full-time workers. The scientifically selected probability sample used was one of many possible samples, each of which could have produced different estimates. A measure of sampling variability for each estimate is available upon request.

SOURCE: Bureau of Labor Statistics, U.S. Department of Labor, February 04, 2010

² Median days away from work is the measure used to summarize the varying lengths of absences from work among the cases with days away from work. Half the cases involved more days and half involved less days than a specified median. Median days away from work are represented in actual values.

Relative standard errors are a measure of the sampling error of an estimate. Sampling errors occur because observations are made on a sample, not on the entire population. Estimates based on the different possible samples of the same size and sample design could differ. Relative standard errors less than 0.05 are not shown.

⁴ Days away from work cases include those which result in days away from work with or without job transfer or restriction.

back; careal tunnel syndrome; hernia; or musculoskeletal system and connective tissue diseases and disorders and when the event or exposure leading to the injury or illness is: bodily reaction/bending, climbing, crawling, reaching, twisting; overexertion; or repetition. Cases of Raynaud's phenomenon, tarsal tunnel syndrome, and herniated spinal discs are not included. Although these cases may be considered MSD's, the survey classifies these cases in categories that also include non-MSD cases.